Neuroscience Letters, 211 (1996) 219–221
Elsevier Science Ireland Ltd.

AUTHOR INDEX

Aikawa, M., see Ueda, S., 97
Akabane, A., see Arai, S., 57
Aksenov, M.Y., see Aksenova, M.V., 45
Aksenova, M.V., Aksenov, M.Y., Butterfield, D.A., Carney, J.M., α-1-Antichymotrypsin interaction with Aβ (1–40) inhibits fibril formation but does not affect the peptide toxicity, 45
Albers, K.M., see Mendelson, B., 68
Andrade, J.P., Paula-Barbosa, M.M., Protein malnutrition alters the cholinergic and GABAergic systems of the hippocampal formation of the adult rat: an immunocytochemical study, 211
Arai, R., see Nagatsu, I., 183
Arai, S., Kinouchi, H., Akabane, A., Owada, Y., Kamii, H., Kawase, M., Yoshimoto, T., Induction of brain-derived neurotrophic factor (BDNF) and the receptor *trk* B mRNA following middle cerebral artery occlusion in rat, 57
Arnaud, M., see Martin-Fardon, R., 179
Aumann, T.D., Horne, M.K., A comparison of the ultrastructure of synapses in the cerebello-rubral and cerebello-thalamic pathways in the rat, 175

Baba, H., see Matsumoto, A., 105
Barman, S.M., see Zhong, S., 113
Barthélémy, P., Sabeur, G., Jammes, Y., Assessment of an airway-to-pulmonary circulation reflex in cats, 89
Benninger, F., see Hoffmann, K., 195
Bertolotto, C., see Liu, R.-H., 163
Bickford, P.C., see Bowenkamp, K.E., 81
Bliss, T.V.P., see Sesay, A.K., 207
Bolten, J., see Hoffmann, K., 195
Bowenkamp, K.E., Lapchak, P.A., Hoffer, B.J., Bickford, P.C., Glial cell line-derived neurotrophic factor reverses motor impairment in 16–17 month old rats, 81
Brent, P.J., Saunders, H., Dunkley, P.R., Intrasynaptosomal free calcium levels in rat forebrain synaptosomes: modulation by sigma (σ) receptor ligands, 138
Butterfield, D.A., see Aksenova, M.V., 45

Cahill, A.L., see Weiss, C., 29
Carney, J.M., see Aksenova, M.V., 45
Challet, E., Malan, A., Pévet, P., Daily hypocaloric feeding entrains circadian rhythms of wheel-running and body temperature in rats kept in constant darkness, 1
Champagnat, J., see Haji, A., 17
Chang, Y.-H., Hsieh, M.-T., Cheng, J.-T., Increase of locomotor activity by acupuncture on *Bai-Hui* point in rats, 121
Chase, M.H., see Liu, R.-H., 163
Chauhan, A., see Wegiel, J., 151
Cheng, J.-T., see Chang, Y.-H., 121
Choi, E.J., see Leem, J.W., 37
Crossman A.R., see Griffiths P.D., 53

Davis, B.M., see Mendelson, B., 68
Dell'Anna, E., Geloso, M.C., Magarelli, M., Molinari, M., Development of GABA and calcium binding proteins immunoreactivity in the rat hippocampus following neonatal anoxia, 93
Deller, T., see Naumann, T., 117
Denavit-Saubié, M., see Haji, A., 17
Dohi, T., see Kitayama, S., 132
Dunkley, P.R., see Brent, P.J., 138

Engelhardt, J.K., see Liu, R.-H., 163
Enomoto, T., see Matsumoto, A., 105
Errington, M.L., see Sesay, A.K., 207

Fetzer, S., see Kress, M., 5
Feucht, M., see Hoffmann, K., 195
Finkelstein, D.I., Reeves, A.K., Horne, M.K., An electron microscopic tracer study of the projections from entopeduncular nucleus to the ventrolateral nucleus of the rat, 33
Fischer-Colbrie, R., see Weiss, C., 29
Foutz, A.S., see Haji, A., 17
Frotscher, M., see Naumann, T., 117
Fujii, J., see Nakano, R., 129
Furuyama, T., see Matsumoto, K., 147

Gebber,Y.,see Zhong, S., 113
Geist, R.T., Gutmann, D.H., Expression of a developmentally-regulated neuron-specific isoform of the neurofibromatosis 1 (NF1) gene, 85
Geloso, M.C., see Dell'Anna, E., 93
Gemba, H., Teranaka, A., Takemura, K., Influences of emotion upon parotid secretion in human, 159
Gericke, C.A., see Riepe, M.W., 9
Goodness, T.P., see Mendelson, B., 68
Granstrom, L., see Zhang, C., 61
Griffiths P.D., Crossman A.R., Autoradiography of transferrin receptors in the human brain, 53
Grönroos, M., Reunala, T., Pertovaara, A., Influence of selective nerve fiber blocks on argon laser-induced thermal pain in the human skin, 143
Gutmann, D.H., see Geist, R.T., 85

Haji, A., Pierrefiche, O., Foutz, A.S., Champagnat, J., Denavit-Saubié, M., Takeda, R., Pharmacological properties of peripherally induced postsynaptic potentials in bulbar respiratory neurons of decerebrate cats, 17
Harada, M., see Mogi, M., 13
Hellweg, R., see Riepe, M.W., 9
Hoffer, B.J., see Bowenkamp, K.E., 81
Hoffmann, K., Feucht, M., Witte, H., Benninger, F., Bolten, J., Analysis and classification of interictal spike discharges in benign partial epilepsy of childhood on the basis of the Hilbert transformation, 195
Horne, M.K., see Aumann, T.D., 175
Horne, M.K., see Finkelstein, D.I., 33
Hsieh, M.-T., see Chang, Y.-H., 121

Iannotti, F., see Pringle, A.K., 203
Inagaki, H., see Mogi, M., 13
Inuzuka, T., see Nakano, R., 129
Ishii, N., see Matsumoto, K., 147
Ishizuya-Oka, A., see Ueda, S., 97

Jammes, Y., see Barthélémy, P., 89
Jibiki, I., see Kubota, T., 21

Kamenka, J.-M., see Martin-Fardon, R., 179
Kamii, H., see Arai, S., 57
Karasawa, N., see Nagatsu, I., 183
Kashihara, Y., see Matsumoto, K., 147
Kasischke, K., see Riepe, M.W., 9
Kauppila, T., Xu, X.-J., Sciatic nerve section induces mechanical hyperalgesia in skin adjacent to the deafferented region in rats: lack of correlation with autotomy behavior, 65
Kawamata, T., see Nakai, M., 41
Kawase, M., see Arai, S., 57
Kawata, M., see Ueda, S., 97
Kikugawa, K., see Nakano, R., 129
Kinouchi, H., see Arai, S., 57
Kishizawa, S., see Kubota, T., 21
Kitanaka, J.-i., see Matsumoto, K., 147
Kitayama, S., Morita, K., Dohi, T., Uptake and release of dopamine through the rat dopamine transporter expressed in *Xenopus laevis* oocyte: evaluation by voltammetric measurement of intracellular dopamine concentration, 132
Kiyatkin, E.A., Stein, E.A., Conditioned changes in nucleus accumbens dopamine signal established by intravenous cocaine in rats, 73
Klein, W.L., see Zhang, C., 187
Krafft, G.A., see Zhang, C., 187
Kream, R.M., see Rittenhouse, P.A., 77
Kress, M., Fetzer, S., Reeh, P.W., Vyklicky, L., Low pH facilitates capsaicin responses in isolated sensory neurons of the rat, 5
Kubota, T., Jibiki, I., Kishizawa, S., Kurokawa, K., Clozapine-induced potentiation of synaptic responses in the perforant path-dentate gyrus pathway in chronically prepared rabbits, 21
Kurokawa, K., see Kubota, T., 21

Lapchak, P.A., see Bowenkamp, K.E., 81
Laslop, A., see Weiss, C., 29
Le Vine, III, H., see Wegiel, J., 151
LeDoux, J.E., see Savander, V., 167
Leem, J.W., Choi, E.J., Park, E.S., Paik, K.S., *N*-Methyl-D-aspartate (NMDA) and non-NMDA glutamate receptor antagonists differentially suppress dorsal horn neuron responses to mechanical stimuli in rats with peripheral nerve injury, 37
Leeman, S.E., see Rittenhouse, P.A., 77
Levita, L., see Sesay, A.K., 207
Li, Y., see Rosenberg, P.A., 49
Liu, G.L., see Zhong, S., 113
Liu, R.-H., Bertolotto, C., Engelhardt, J.K., Chase, M.H., Age-related changes in soma size of neurons in the spinal cord motor column of the cat, 163
Löwe, A., see Riepe, M.W., 9

Maeda, K., see Nakai, M., 41
Magarelli, M., see Dell'Anna, E., 93
Malan, A., see Challet, E., 1
Marchand, J.E., see Rittenhouse, P.A., 77
Markle, R.A., see Rittenhouse, P.A., 77
Martin-Fardon, R., Arnaud, M., Rousseau, É., Kamenka, J.-M., Privat, A., Vignon, J., *N*-[1-(2-Benzo(*b*)thiophenyl)cyclo-hexyl]piperidine (BTCP) and cocaine induce similar effects on striatal dopamine: a microdialysis study in freely moving rats, 179
Maruyama, K., see Nishimura, E., 101
Matsumoto, A., Matsumoto, R., Enomoto, T., Baba, H., Human brain β-secretase contains heparan sulfate glycoconjugates, 105
Matsumoto, K., Tanaka, T., Furuyama, T., Kashihara, Y., Ishii, N., Tohyama, M., Kitanaka, J.-i., Takemura, M., Mori, T., Wanaka, A., Differential expression of LIM-homeodomain genes in the embryonic murine brain, 147
Matsumoto, R., see Matsumoto, A., 105
Mendelson, B., Albers, K.M., Goodness, T.P., Davis, B.M., Overexpression of nerve growth factor in epidermis of transgenic mice preserves excess sensory neurons but does not alter the somatotopic organization of cutaneous nerve projections, 68
Minami, M., see Mogi, M., 13
Mogi, M., Harada, M., Narabayashi, H., Inagaki, H., Minami, M., Nagatsu, T., Interleukin (IL)-1β, IL-2, IL-4, IL-6 and transforming growth factor-α levels are elevated in ventricular cerebrospinal fluid in juvenile parkinsonism and Parkinson's disease, 13
Molinari, M., see Dell'Anna, E., 93
Mori, T., see Matsumoto, K., 147
Morita, K., see Kitayama, S., 132

Nagatsu, I., Takeuchi, T., Sakai, M., Karasawa, N., Yamawaki, Y., Arai, R., Nagatsu, T., Transient appearance of tyrosine hydroxylase-immunoreactive non-catecholaminergic neurons in the medial geniculate nucleus of postnatal mice, 183
Nagatsu, T., see Mogi, M., 13
Nagatsu, T., see Nagatsu, I., 183
Nakai, M., Kawamata, T., Taniguchi, T., Maeda, K., Tanaka, C., Expression of apolipoprotein E mRNA in rat microglia, 41
Nakano, R., Inuzuka, T., Kikugawa, K., Takahashi, H., Sakimura, K., Fujii, J., Taniguchi, N., Tsuji, S., Instability of mutant Cu/Zn superoxide dismutase (Ala4Thr) associated with familial amyotrophic lateral sclerosis, 129
Narabayashi, H., see Mogi, M., 13
Naumann, T., Deller, T., Frotscher, M., Multiple projections are unlikely to account for the survival of rat medial septal neurons after axotomy, 117
Nichols, R.A., Suplick, G.R., Rapid chelation of calcium entering isolated rat brain nerve terminals during stimulation inhibits neurotransmitter release, 135
Nishimura, A., see Ueda, S., 97
Nishimura, E., Sasaki, K., Maruyama, K., Tsukada, T., Yamaguchi, K., Decrease in neuron-restrictive silencer factor (NRSF) mRNA levels during differentiation of cultured neuroblastoma cells, 101
Nowakowski, J., see Wegiel, J., 151

Obata, K., see Yamagata, Y., 109
Obata, T., Yamanaka, Y., Cardiac microdialysis of salicylic acid to detect hydroxyl radical generation associated with sympathetic nerve stimulation, 216
Okada, Y.C., Xu, C., Single-epoch neuromagnetic signals during epileptiform activities in guinea pig longitudinal CA3 slices, 155
Owada, Y., see Arai, S., 57

Paik, K.S., see Leem, J.W., 37
Park, E.S., see Leem, J.W., 37
Paula-Barbosa, M.M., see Andrade, J.P., 211
Perlman, R.L., see Weiss, C., 29
Pertovaara, A., see Grönroos, M., 143
Pévet, P., see Challet, E., 1
Pierrefiche, O., see Haji, A., 17
Pitkänen, A., see Savander, V., 167

Pringle, A.K., Sundstrom, L.E., Wilde, G.J.C., Williams, L.R., Iannotti, F., Brain-derived neurotrophic factor, but not neurotrophin-3, prevents ischaemia-induced neuronal cell death in organotypic rat hippocampal slice cultures, 203
Privat, A., see Martin-Fardon, R., 179

Qiu, H.E., see Zhang, C., 187

Reeh, P.W., see Kress, M., 5
Reeves, A.K., see Finkelstein, D.I., 33
Reunala, T., see Grönroos, M., 143
Riepe, M.W., Kasischke, K., Gericke, C.A., Löwe, A., Hellweg, R., Increase of hypoxic tolerance in rat hippocampal slices following 3-nitropropionic acid is not mediated by endogenous nerve growth factor, 9
Rittenhouse, P.A., Markle, R.A., Marchand, J.E., Kream, R.M., Leeman, S.E., Streptozotocin-induced diabetes produces a decrease in pituitary substance P content and preprotachykinin mRNA, 77
Rosenberg, P.A., Li, Y., Forskolin evokes extracellular adenosine accumulation in rat cortical cultures, 49
Rossiter, C.D., Yates, B.J., Vestibular influences on hypoglossal nerve activity in the cat, 25
Rousseau, É., see Martin-Fardon, R., 179

Sabeur, G., see Barthélémy, P., 89
Sakai, M., see Nagatsu, I., 183
Sakimura, K., see Nakano, R., 129
Sasaki, K., see Nishimura, E., 101
Saunders, H., see Brent, P.J., 138
Savander, V., LeDoux, J.E., Pitkänen, A., Topographic projections from the periamygdaloid cortex to select subregions of the lateral nucleus of the amygdala in the rat, 167
Schultz, K., Stell, W.K., Immunocytochemical localization of the high-affinity glutamate transporter, EAAC1, in the retina of representative vertebrate species, 191
Sesay, A.K., Errington, M.L., Levita, L., Bliss, T.V.P., Spatial learning and hippocampal long-term potentiation are not impaired in *mdx* mice, 207
Sheetz, M.P., see Toyoshima, I., 171
Shirasawa, T., see Yahagi, Y.-i., 125
Starkey, S.J., Melatonin and 5-hydroxytryptamine phase-advance the rat circadian clock by activation of nitric oxide synthesis, 199
Stein, E.A., see Kiyatkin, E.A., 73
Stell, W.K., see Schultz, K., 191
Sundstrom, L.E., see Pringle, A.K., 203
Suplick, G.R., see Nichols, R.A., 135

Tagawa, M., see Yahagi, Y.-i., 125
Takahashi, H., see Nakano, R., 129
Takeda, R., see Haji, A., 17
Takemura, K., see Gemba, H., 159
Takemura, M., see Matsumoto, K., 147
Takeuchi, T., see Nagatsu, I., 183
Tanaka, C., see Nakai, M., 41
Tanaka, T., see Matsumoto, K., 147
Taniguchi, N., see Nakano, R., 129
Taniguchi, T., see Nakai, M., 41
Teranaka, A., see Gemba, H., 159
Tohyama, M., see Matsumoto, K., 147
Tomoda, T., see Yahagi, Y.-i., 125
Toyoshima, I., Sheetz, M.P., Kinectin distribution in chicken nervous system, 171
Tsuji, S., see Nakano, R., 129
Tsukada, T., see Nishimura, E., 101

Ueda, S., Aikawa, M., Ishizuya-Oka, A., Nishimura, A., Kawata, M., Alteration of serotonergic innervation in the suprachiasmatic nucleus of the rat following removal of input fibers from retina and lateral geniculate nucleus, 97

Vignon, J., see Martin-Fardon, R., 179
Vyklicky, L., see Kress, M., 5

Wanaka, A., see Matsumoto, K., 147
Wang, K.-C., see Wegiel, J., 151
Wegiel, J., Chauhan, A., Wisniewski, H.M., Nowakowski, J., Wang, K.-C., Le Vine, III, H., Promotion of synthetic amyloid β-peptide fibrillization by cell culture media and cessation of fibrillization by serum, 151
Weiss, C., Cahill, A.L., Laslop, A., Fischer-Colbrie, R., Perlman, R.L., Winkler, H., Differences in the composition of chromaffin granules in adrenaline and noradrenaline containing cells of bovine adrenal medulla, 29
Wilde, G.J.C., see Pringle, A.K., 203
Williams, L.R., see Pringle, A.K., 203
Winkler, H., see Weiss, C., 29
Wisniewski, H.M., see Wegiel, J., 151
Witte, H., see Hoffmann, K., 195
Wong-Riley, M.T.T., see Zhang, C., 61

Xu, C., see Okada, Y.C., 155
Xu, X.-J., see Kauppila, T., 65

Yahagi, Y.-i., Tagawa, M., Tomoda, T., Shirasawa, T., Binary expression of olfactory bulb-protein tyrosine phosphatase in rat central nervous system: developmental gene regulation in neonate cerebral cortex and constitutive expression in olfactory-rhinencephalon, 125
Yamagata, Y., Obata, K., Ca^{2+}/calmodulin-dependent protein kinase II in septally kindled rat brains: changes in protein level, activity and subcellular distribution in hippocampus and cerebral cortex, 109
Yamaguchi, K., see Nishimura, E., 101
Yamanaka, Y., see Obata, T., 216
Yamawaki, Y., see Nagatsu, I., 183
Yates, B.J., see Rossiter, C.D., 25
Yoshimoto, T., see Arai, S., 57

Zhang, C., Granstrom, L., Wong-Riley, M.T.T., Deafferentation leads to a down-regulation of nitric oxide synthase in the rat visual system, 61
Zhang, C., Qiu, H.E., Krafft, G.A., Klein, W.L., Aβ peptide enhances focal adhesion kinase/Fyn association in a rat CNS nerve cell line, 187
Zhong, S., Gebber,Y., Liu, G.L., Zhou, S.-Y., Barman, S.M., Comparison of results obtained with time- and frequency-domain analysis when searching for the 10-Hz rhythm in sympathetic nerve discharge, 113
Zhou, S.-Y., see Zhong, S., 113